Suggested Duration of Therapy for Common Infectious Diseases*

Site of Infection	Clinical Diagnosis	Duration of Therapy
Blood	Bacteremia	
	- Staphylococcus aureus	Minimum 14 days from 1st negative culture
	 Uncomplicated Gram-negative 	7 days
	 Complicated Gram-negative 	14 days
	 Uncomplicated CoNS catheter- 	5-7 days
	related (with line removal)	
	 All other organisms 	14 days from 1 st negative culture
Heart	Infective endocarditis	
	 Viridans Group Strep 	2-6 weeks
	- Enterococcus spp.	4-6 weeks
	- Staphylococcus aureus	6 weeks
	Meningitis	
	- Neisseria meningitidis	7 days
	- Haemophilus influenzae	7 days
Brain	- Streptococcus pneumoniae	10 days
	- Listeria monocytogenes	21 days
	- Gram-negative	14-21 days
	- HSV or VZV	21 days
Bone	Osteomyelitis	6 weeks
Joint	Septic arthritis	14 days
	Gonococcal arthritis	7 days
	Prosthetic joint infection	,
	- Retention	6 weeks (non-Staph), 3-6 months (Staph)
	- One-step exchange	6 weeks (non-Staph), 3 months (Staph)
	- Two-step exchange	6 weeks
	- Permanent resection	6 weeks
	- Amputation	24 hours if adequate surgical margins
GI	Cholecystitis	4-7 days
	Acute uncomplicated diverticulitis	4-7 days
	Acute uncomplicated appendicitis	4-7 days

	Peritonitis	4-7 days
	Clostridioides difficile infection	10 days
Genitourinary	Asymptomatic bacteriuria	Do not treat ^
	Cystitis	3-5 days
	Pyelonephritis	7 days
	Acute Bacterial Prostatitis	4 weeks
	Chronic Bacterial Prostatitis	4-6 weeks
Lung	CAP	5 days
	HAP/VAP	7 days
	Legionella pneumonia	7 days
	PJP pneumonia	21 days
Skin	Cellulitis	5 days
	Purulent SSTI (with I&D)	Outpatient: 5 days, Inpatient: 7 days
	Bite Wound	7 days
	Diabetic Foot Infection (no osteomyelitis)	Mild or Moderate: 7 days, Severe: 14 days
	Necrotizing fasciitis	7 days after last surgical debridement
	Surgical site infection	24 hours after surgical debridement

^{*}Assuming clinical improvement
^Exceptions: pregnant, renal transplant within past 1 month, undergoing traumatic urologic procedure