For more details. **NM West Region Allergy Algorithm** [†]Be sure to document in click here to go to patient chart if an System Allergy Risk alternative beta-lactam Reported Beta-Lactam Reaction[†] Assessment is tolerated Severe, non-IgE-Severe, Nonmediated IgE-Severe mediated (e.g., Stevens-Johnson syndrome, TEN, serum sickness, DRESS, liver or kidney damage) History IgE-Like **Anaphylaxis** inconsistent with Unknown Reaction* ≥10 Or allergy Or years ago Non-Severe IgE-Like Reaction* <10 (e.g., rash, (e.g., side effect or **EXCLUDING** family history but hives) years ago **Anaphylaxis** no patient reaction) Give alternate/ non-beta-Cephalo-Cephalo-**PCN** PCN lactam sporin sporin **Allergy Allergy** antibiotic **Allergy** Allergy *IgE-Like Reaction: Cefazolin can be A PCN or a Cutaneous: Itching, flushing, hives, angioedema A cephalosporin Give **Discuss with** safely administered cephalosporin Respiratory: Shortness of breath, rhinitis, provider to consider cephalosporin with different alternate/ (if indicated) wheezing, bronchospasm with different delabeling allergy can be safely side-chain can non-beta-GI symptoms: Immediate nausea, vomiting, side-chain can diarrhea, abdominal pain (usually related to Otherwise, give administered lactam be safely be safely alternate/non-betaabove symptoms, not as in a side effect) antibiotic administered lactam antibiotic administered